

Accepted Manuscript

BJGP OPEN

Clinical and serological profile of asymptomatic and non-severe symptomatic COVID-19 cases: Lessons from primary care in Latin America

Klaus Puschel, Catterina Ferreccio, Blanca Peñaloza, Katia Abarca, Maria-Paulina Rojas, Alvaro Tellez, Philippa Moore, Ana Maria Cea, | Carlos Wilson, Vicente Cid, Joaquin Montero

DOI: <https://doi.org/10.3399/bjgpopen20X101137>

To access the most recent version of this article, please click the DOI URL in the line above.

Received 17 August 2020

Revised 07 September 2020

Accepted 10 September 2020

© 2020 The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>). Published by BJGP Open. For editorial process and policies, see: <https://bjgpopen.org/authors/bjgp-open-editorial-process-and-policies>

When citing this article please include the DOI provided above.

Author Accepted Manuscript

This is an 'author accepted manuscript': a manuscript that has been accepted for publication in BJGP Open, but which has not yet undergone subediting, typesetting, or correction. Errors discovered and corrected during this process may materially alter the content of this manuscript, and the latest published version (the Version of Record) should be used in preference to any preceding versions

Clinical and serological profile of asymptomatic and non-severe symptomatic Covid-19 cases: Lessons from primary care in Latin America.

Puschel, Klaus; Ferreccio, Catterina; Peñaloza, Blanca; Abarca, Katia; Rojas, Maria-Paulina; Tellez, Alvaro; Moore, Philippa; Cea, Ana Maria; Wilson, Carlos; Cid, Vicente; Montero, Joaquin

Klaus Puschel MD, MPH, MSc
Professor.

Department of Family and Community Medicine¹.
School of Medicine, Pontificia Universidad Católica de Chile ²
Av. Diagonal Paraguay 362, Edificio Académico Escuela de Medicina UC, Santiago, Chile³
kpuschel@med.puc.cl

Catterina Ferreccio MD, MPH ^{2,3}
Professor
Department of Public Health⁴.
Epidemiology PhD Program Director.

Blanca Peñaloza MD, MSc ^{1,2,3}
Associate Professor

Katia Abarca MD, MSc ^{2,3}
Professor
Department of Pediatric Infectious Diseases.

Maria Paulina Rojas MD ^{1,2,3}
Associate Professor
Chair, Department of Family and Community Medicine.

Alvaro Tellez MD, MPH ^{1,2,3}
Associate Professor

Philippa Moore MD, MSc ^{1,2,3}
Associate Professor

Ana Maria Cea RN ^{1,2,3}
Clinical Research Assistant

Carlos Wilson MD ^{1,2,3}
Clinical Research Assistant

Vicente Cid BSc ^{2,3,4}
Statistical Research Assistant

Joaquin Montero MD, MPH ^{1,2,3}
Professor

Clinical and serological profile of asymptomatic and non-severe symptomatic Covid-19 cases: Lessons from primary care in Latin America.

Key words: Covid-19, primary health care, Latin America

How this fits in:

- Chile has one of the highest incidence of Covid-19 in the world. Primary care can play a key role in early detection of cases and containment of the epidemic.
- The clinical and serological profile of Covid-19 has not been thoroughly evaluated in the large population of asymptomatic or non-severe symptomatic cases seen in primary care.
- The immune response of asymptomatic and non-severe cases is significantly lower than that reported in severe cases. Symptoms profile also differs in both populations.
- Rapid serologic testing is ineffective for detecting asymptomatic or non-severe symptomatic cases of Covid-19 but can be of value for surveillance of immunity response at the primary care and community levels.

Abstract:

Background: Chile, presents one of the highest incidences of Covid-19 infection in the world. Primary care can play a key role in early detection and contention of the disease. There is lack of information on the clinical profile of suspected Covid-19 primary care patients and controversy on the effectiveness of rapid serologic tests in the diagnosis and surveillance of the disease.

Aim: To assess the effectiveness of rapid serologic testing in detection and surveillance of Covid-19 cases in primary care.

Design and Setting: Longitudinal study based on non-random sample of 522 participants including 304 symptomatic patients and 218 high-risk asymptomatic individuals receiving care at four primary health clinics in an underserved area in Santiago.

Method: Participants were systematically assessed and tested for Covid-19 with RT-PCR and serology at baseline and followed clinically and serologically for a three weeks period.

Results: The prevalence rate of RT-PCR confirmed Covid-19 cases was 3.5 times higher in symptomatic patients (27.5% (22.1-32.8; 95% CI) compared to asymptomatic participants (7.9% (4.3-11.6; 95% CI). Similarly, the immune response was significantly different between both groups. Sensitivity of serologic testing was 57.8% (44.8-70.1; 95% CI) during the third week of follow-up and specificity was 98.4% (95.5-99.7; 95% CI).

Conclusion: Rapid serologic testing is ineffective for detecting asymptomatic or non-severe cases of COVID-19 at early stages of the disease but can be of value for surveillance of immunity response in primary care. The clinical profile and immune response of Covid-19 patients in primary care differ from those in hospital-based populations.

Introduction

The pandemic of Covid-19 is emerging strongly in Latin America, by mid-July the number of new daily cases surpassed that of the United States or Europe with Brazil, Perú and Chile leading the incidence rates. Chile presented one of the highest prevalence rate of Covid-19 in the world; higher than Brazil, the United States or the United Kingdom as of August 31st, 2020 (1). The recent stabilization of incidence rates reported in Chile and in other countries highlights the relevance of finding effective early diagnostic and surveillance strategies at the community level to contain the epidemic and start a safe unlock process.

Primary Health Care is at the frontline of the epidemic interface between the community and health care services (2). Over 80% of Covid-19 cases are non-severe cases that remain in the community and are detected and surveilled in primary care (3). The clinical and serological profile of asymptomatic or non-severe cases that present with mild symptoms has not been addressed thoroughly. The great majority of studies evaluating the clinical and serological profile of the Covid-19 infected population have been conducted in hospitalized patients with severe disease (4–6). Consistent information on the clinical and serological profile of non-severe Covid-19 cases could contribute to early diagnosis, contact tracing and effective isolation strategies. As lockdown strategies are starting to be reduced, due to a decrease in cases or due to social exhaustion, the role of primary care services could be essential for effective detection and containment of new outbreaks.

Effective testing and tracing of Covid-19 cases is a key factor in the successful strategies for controlling the epidemic at the primary care level. Accuracy, acceptability, affordability and results availability of the test at the point of care are essential requirements for effective testing (7). The standard reverse transcriptase-polymerase chain reaction test (RT-PCR) testing has important limitations as a massive test strategy at the primary care level given that it requires a special sampling procedure, central lab processing and has a latency of 24-72 hours for the results. Latency time of the test results can be especially problematic in an epidemic scenario where instant results are needed to implement immediate isolation measures. In addition, sensitivity of RT-PCR declines significantly after 7-8 days of infection with reported false negative rates of 30-40% (8).

Rapid serologic tests have the potential to be an alternative to face the challenge of detection and tracing cases in highly demanded primary care clinics in an epidemic scenario. However, high variability in the accuracy of serologic tests have been reported. In a recent meta-analysis of 38 studies that included 7848 participants, Kontou P et al (9), found a wide sensitivity rates that varied from 56% to 83% and specificity rates that ranged from 91% to 99% depending on the characteristics of the serologic test used and the population studied. Many studies included in the meta-analysis did not report clinical severity and were underpowered. Similar results were found in a recent meta-analysis conducted by Lisboa-Bastos M et al (10) that analyzed 40 studies but only two of them included outpatients and point of care testing. These results do not permit decision makers to draw consistent conclusions on the real value of serologic testing in primary care.

This study assessed the demographic, clinical and serological profile of non-severe symptomatic and asymptomatic cases of Covid-19 in primary care setting. The study

evaluated the accuracy of rapid serologic testing in the detection and follow-up of COVID-19 infections in primary care

Methods:

Population and sampling

The study was conducted in four primary health care clinics at the South East area of Santiago, Chile. This area has an estimated population of 1 million people of middle and low socioeconomic status. A non-random sample of 522 individuals was included in the study, 304 of these were symptomatic adults who consulted at these clinics between April 8th and May 14th-2020, and fulfilled the criteria defined by the World Health Organization of suspected cases or had any respiratory symptoms at the time of the visit. In addition, 218 asymptomatic individuals belonging to high risk groups such as close contacts of confirmed cases, primary care health personnel or prison officers working in high risk environments in the study area were also invited to participate the study.

Procedures

Patients who accepted to participate in the study were assessed clinically using the national clinical guideline recommended for suspected cases (11). The guideline included assessing for demographic and risk factors (e.g. smoking, high blood pressure, diabetes), chronic diseases (e.g. asthma, COPD, cardiovascular disease), medications (e.g. ARBs, NSAID) respiratory (e.g. sore throat, cough, fever, dyspnea) and other symptoms (myalgia, headache, anosmia, dysgeusia, gastrointestinal symptoms). It also included a physical exam and general laboratory testing or imaging as required.

At entry, (Day 1) all participants were tested for Covid-19 through a standard RT-PCR test using a nasopharyngeal swab. The samples were processed at the Microbiology Laboratory, UC-Christus Health Network.. Additionally, participants were tested with the rapid serologic test. The test was design to identify serum IgM and IgG from a finger capillary blood that was taken using a point-of-care (POC) Lateral Flow Immunoassays (LFIA) Covid-19 test (Acro-Biotec Inc, CA 91730, USA). Results were obtained 10 minutes after the finger stick. Participants were re-evaluated at Day 7 if they tested PCR negative at entry and all were re-tested and a received a brief clinical assessment after Day 14.

The study was approved by the Ethical Committee of Health Sciences at Pontificia Universidad Católica de Chile and also by the local Committee at the South East Health Services Area in Santiago, Chile. All patients read and signed an informed consent before entering the study. The study was funded by Pontificia Universidad Católica de Chile and by BHP Billiton mining company, Melbourne, Australia as a part of the Esperanza Covid-19 Project (12).

Analysis

We conducted stratified analysis of symptomatic and asymptomatic participants, and estimated sensitivity, specificity and likelihood ratios for serologic tests at baseline and follow-up. We explored associations between risk factors, symptoms, and RT-PCR with serologic results with multivariate logistic regression using R 3.5.1 and Stata 15.

Results

The four groups included in the study differed in demographic characteristics such as age, gender and also in the presence of chronic diseases (Table 1). Symptomatic patients were older (42.5 and 33.6 years, $p < 0.001$), used more medications (28.1% and 10.5%, $p < 0.001$) and had a higher history of chronic respiratory diseases (8.7% and 1.8%, $p < 0.05$) compared to all asymptomatic participants. They were similar regarding prevalence of smoking, high blood pressure and diabetes. The median interval between the first symptom and the clinical assessment in the symptomatic group was three days and the most frequent symptoms were headache, dry cough and sore throat. The positivity of RT-PCR test was 3.5 times higher in symptomatic vs. asymptomatic groups (27.5% (22.1-32.8) vs. 7.9% (4.3-11.6); $p < 0.001$). Similarly, positivity of IgM or IgG during the study period among symptomatic patients was 3.8 times higher than the asymptomatic participants (18.5% (14.1-23.0) vs. 4.9 (1.7-8.1); $p < 0.001$).

The relative sensitivity, specificity and likelihood ratios of the serologic tests compared to RT-PCR in symptomatic patients over time are shown in Table 2. The relative sensitivity of serology increased from 4.9% in the first week to 57.8% at Day 14. The specificity of serological tests remained over 96% throughout the study. Positive likelihood ratio was 36.6 after 14 days of initial symptoms. The sensitivity of the serologic tests in asymptomatic participants remained very low, below 5% during the follow-up period and specificity remained over 97%. Figure 1 presents the positivity rate of serologic tests in symptomatic and asymptomatic patients with positive RT-PCR results over time. Main predictors of a positive serology among symptomatic patients were: older age (PR: 1.03 (1.02-1.05) , female gender (PR: 1.63 (1.06-2.51)) , and dry cough (PR: 1.79 (1.14-2.81)).

Discussion

Summary

This study shows that LFIA serologic testing is not effective for early diagnosis of Covid-19 in primary health care given its low accuracy in detecting cases. The immune response of asymptomatic and non-severe cases is significantly lower than that reported in severe cases. The symptom profile also differs in both populations. LFIA serologic testing might be of value for surveillance in primary health care and in the community given its high specificity and increasing detection rate after two weeks of infection.

Strengths and limitations

This study targets a key population for controlling the epidemic that has not been addressed thoroughly. Participants were non-randomly selected and therefore do not represent an homogenous population. They were selected on clinical criteria of symptoms suggestive of Covid-19 disease or on epidemiological criteria of high-risk populations. The criteria used for including participants were similar to that recommended for Covid-19 testing in primary health care and in the community (13) and our study reflects the reality of this level of care. The LFIA tests used in this study had the least sensitivity of the serologic tests available for rapid testing. The test was chosen because it was the most widely used at the time, was affordable, rapidly available and the reported accuracy was acceptable. These are essential characteristics to value the effectiveness of a diagnostic test in primary care (7).

Comparison with existing literature

The sero-prevalence results of our study are similar than those observed in a community-based study conducted in England that reported 7% of sero-prevalence in May 2020 (14), and those reported in the surveillance program in England which found a range from 4% (South East) to 16% (London) at week 2020 in a mixed adult population (15). In contrast to severe Covid-19 cases where fever is the predominant symptom present in over 80% of cases (4), in our non-severe population fever was present in only 15 % of cases. Dry cough was the predominant symptom in our symptomatic population with a significant predictive value for infection.

Accuracy assessment of serologic testing in our study revealed a low relative sensitivity that ranged from 8% during the first week of symptoms or contact date to 58% during the third week of follow-up. Specificity remained over 95% along the study when compared to RT-PCR results. The low sensitivity observed in this study contrasts with most reports that showed sensitivity rates over 80% (9) and also with the manufactured company data of the serologic POC test used in this study that reported relative sensitivity values of 100% for IgM and 85% for IgG (16). These results highlight the difference in the intensity of humoral response in non-severe cases versus in the severe cases that have also been reported in other studies (17,18).

The results of this investigation are consistent with evidence showing that there is a significant seroconversion rate of IgG after 14 days of symptom onset but, that antibody levels in non-severe cases are low or maybe undetectable (17,19,20). The increase in the seroconversion rate over time observed was associated, as expected, with a raise in the sensitivity of the test after the second week that has also been observed in many other studies (10). The available evidence shows that viral dynamics and serologic response are very different when comparing asymptomatic, non-severe or severe Covid-19 cases. Liu et al (21) found that the mean viral load of severe cases was around 60 times higher than that detected in mild cases in a group of 76 patients admitted to the Nanchang University Hospital in China. Similarly, Long et al (5) found that severely ill patients had significantly higher IgG levels compared with non-severe patients in a cohort of 250 hospitalized patients in Hubei, China. Our results are also consistent with evidence showing a very low seroconversion in

asymptomatic versus symptomatic patients and high percentage of seronegative conversion after the early convalescent phase (17,18). Our study shows a relatively low detectable seroconversion rate in about 60% of our non-severe population after the second week of infection with a significant difference between symptomatic and asymptomatic high-risk patients. By providing specific information in a population of non-severe and asymptomatic cases, this study contributes to the growing evidence of a gradient between severity of the disease, viral load and immune response.

The characteristics of the test used for antibody detection is another important source of variability in the detection rate of Covid-19 infection. In a meta-analysis of 37 studies conducted by Kountou et al (9), specificity of the four techniques analyzed were over 90%. Our results are consistent with these estimates. However, the sensitivity of the tests was much more variable and lateral flow immunoassays (LFIA) were the least sensitive with a range of 75%-80%. The National COVID Advisory Panel at the UK, found that sensitivity estimates for LFIA ranged from 55-70% versus 65-85% for Enzyme-linked immunosorbent assays (ELISA) when compared to RT-PCR (22). However, most of the studies analyzed by both groups included only hospitalized and severe patients. Our study found a sensitivity of 57.8% at the third week of infection in our population of non-severe symptomatic and asymptomatic cases.

Implications for Research and/or practice

Early detection of non-severe and asymptomatic Covid-19 cases has been defined as a key strategy to control the epidemic (15) given that these cases account for about 80% of the infected population (3). Our study targets this population and shows that given its low sensitivity, LFIA tests are not effective to detect Covid-19 in early stages in non-severe symptomatic or high-risk asymptomatic populations. As such, LFIA appear to be of little use in the early diagnosis of COVID-19 at a primary care level. Other techniques such as ELISA, or RT-PCR saliva test might be more effective techniques for early diagnosis but at a higher cost.

In spite of the limitations of the serologic testing observed in this study for early diagnosis in a population with non-severe Covid-19 infection, LFIA could be of value for surveillance at a community level given the high specificity observed after the second week of infection. Applying the rapid test in a closed community such as a school, workplace or neighborhood could help to estimate the prevalence of past disease, magnitude of recent outbreak and the real need for new testing. The presence of detectable levels of antibodies mean that infection have already occurred with a very high probability, even in individuals with low intensity of symptoms. It can provide an estimate of the immune status of a local community (23). The experience in Singapore (24) and in Taiwan (25) showed that serologic testing is a complementary strategy for surveillance given that RT-PCR fails to detect cases beyond the first week of infection. Serologic testing can provide valuable information on the attack and transmission rates at the community level (23). These are key factors to consider for improving containment of an outbreak or defining unlocking measures at a local level (14). Our study shows an increasing serologic trend beyond the second week of infection and a very low false positive rates in symptomatic and asymptomatic cases. These results can contribute to improve surveillance of the disease at the community level.

References

1. Johns Hopkins Coronavirus Resource Center. 2020
<https://coronavirus.jhu.edu/map.html> (accessed 31 August 2020)
2. Kidd MR. Five principles for pandemic preparedness: lessons from the Australian COVID-19 primary care response. *Br J Gen Pract* 2020; 70 (696):316-317
3. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China. *JAMA* 2020;323(13):1239-1242:
4. Liang W, Liang H, Ou L, et al. Development and Validation of a Clinical Risk Score to Predict the Occurrence of Critical Illness in Hospitalized Patients With COVID-19. *JAMA Intern Med* 2020 E1-E9
5. Long Q-X, Liu B-Z, Deng H-J, et al. Antibody responses to SARS-CoV-2 in patients with COVID-19. *Nat Med* 2020; 26(6):845-848
6. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395(10229):1054–1062.
7. Kosack CS, Page A-L, Klatser PR. A guide to aid the selection of diagnostic tests. *Bull World Health Organ* 2017;95(9):639–645.
8. Kucirka LM, Lauer SA, Laeyendecker O, et al. Variation in False-Negative Rate of Reverse Transcriptase Polymerase Chain Reaction–Based SARS-CoV-2 Tests by Time Since Exposure. *Ann Intern Med* 2020;M20-1495.
9. Kontou PI, Braliou GG, Dimou NL, et al. Antibody tests in detecting SARS-CoV-2 infection: a meta-analysis. *medRxiv* 2020 Apr 25
<https://www.medrxiv.org/content/10.1101/2020.04.22.20074914v1> (accessed 12 July 2020)
10. Lisboa Bastos M, Tavaziva G, Abidi SK, et al. Diagnostic accuracy of serological tests for covid-19: systematic review and meta-analysis. *BMJ* 2020;370:m2516.
11. Colegio Médico. Recomendaciones sobre manejo del COVID 19 en Chile. Coordinación Sociedades Científicas y Colegio Médico de Chile. Santiago; 2020
http://www.colegiomedico.cl/wp-content/uploads/2020/05/Declaración-Estrategia-Sanitaria-Sociedades-y-COLMED-29_05-1.pdf (accessed 5 July 2020)
12. Chile PUC de. UC y BHP desarrollan estrategia para mejorar la respuesta de Chile frente a Covid-19: Proyecto Esperanza Covid-19. Portal Noticias UC 2020;
<https://www.uc.cl/noticias/uc-y-bhp-desarrollan-estrategia-para-mejorar-la-respuesta-de-chile-frente-a-covid-19/> (accessed 2 July 2020)

13. MacIntyre CR. Case isolation, contact tracing, and physical distancing are pillars of COVID-19 pandemic control, not optional choices. *Lancet Infect Dis* 2020;S1473-3099(20)30512-0
14. Baraniuk C. Covid-19 antibody tests: a briefing. *BMJ* 2020;369:m2284.
15. Public Health England. Sero-surveillance of COVID-19 - GOV.UK 2020. <https://www.gov.uk/government/publications/national-covid-19-surveillance-reports/sero-surveillance-of-covid-19> (accessed 11 July 2020)
16. Acro Biotech Inc. 2019-nCov IgG/IgM Rapid Cassette. California, U.S.A. [https://www.assaygenie.com/content/Acro Biotech/ACRO Pack Insert.pdf](https://www.assaygenie.com/content/Acro%20Biotech/ACRO%20Pack%20Insert.pdf) (accessed 6 June 2020)
17. Long Q-X, Tang X-J, Shi Q-L, et al. Clinical and immunological assessment of asymptomatic SARS-CoV-2 infections. *Nat Med* 2020 <http://www.nature.com/articles/s41591-020-0965-6> (accessed 7 July 2020)
18. Yongchen Z, Shen H, Wang X, et al. Different longitudinal patterns of nucleic acid and serology testing results based on disease severity of COVID-19 patients. *Emerg Microbes Infect* 2020;9(1):833–6.
19. Kellam P, Barclay W. The dynamics of humoral immune responses following SARS-CoV-2 infection and the potential for reinfection. *J Gen Virol* 2020 <https://www.microbiologyresearch.org/content/journal/jgv/10.1099/jgv.0.001439> (accessed 14 June 2020)
20. Zhao J, Yuan Q, Wang H, et al. Antibody responses to SARS-CoV-2 in patients of novel coronavirus disease 2019. *Clin Infect Dis* 2020 <https://academic.oup.com/cid/advance-article/doi/10.1093/cid/ciaa344/5812996> (accessed 2 July 2020)
21. Liu Y, Yan L-M, Wan L, et al. Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis* 2020;20(6):656-657.
22. Adams ER, Ainsworth M, Anand R, et al. Antibody testing for COVID-19: A report from the National COVID Scientific Advisory Panel. *medRxiv* 2020 <https://www.medrxiv.org/content/10.1101/2020.04.15.20066407v2> (accessed 15 June 2020)
23. Winter AK, Hegde ST. The important role of serology for COVID-19 control. *Lancet Infect Dis* 2020 20(7):758-759
24. Yong SEF, Anderson DE, Wei WE, et al. Connecting clusters of COVID-19: an epidemiological and serological investigation. *Lancet Infect Dis* 2020; 20 (7): 809–815
25. Wang CJ, Ng CY, Brook RH. Response to COVID-19 in Taiwan: Big Data Analytics, New Technology, and Proactive Testing. *JAMA* 2020; 323(14):1341-1342

Accepted Manuscript – BJGP Open – bjgpopen20X101137

Accepted Manuscript – BJGP Open – bjgpopen20X101137

Table 1. Characteristics of study participants (n=522)

	Symptomatic patients (n=304, 58.2%)	Asymptomatic patients (n= 218, 41.8%)		
		Prison Officers (n= 139, 26.6%)	Health Workers (n=39, 7.5%)	Contacts (n=40, 7.7%)
Charateristics	Mean / Prevalence % (95% CI)	Mean / Prevalence % (95% CI)	Mean / Prevalence % (95% CI)	Mean / Prevalence % (95% CI)
Age (yrs) mean ± sd	42.5 ± 15.3	32.7 ± 6.8	33.0 ± 9.0	37.3 ± 13.8
Women (n=217)	53.6 (48.0-59.2)	5.8 (1.9-9.6)	76.9 (63.7-90.1)	40.0 (24.8-55.2)
Current smoker (n=149)	34.8 (28.9-40.8)	37.3 (27.5-47.2)	22.6 (6.7-38.5)	27.5 (12.3-42.7)
Medications:				
ARBs (n=37)	9.0 (6.1-11.9)	-	10.1 (0.0-21.6)	17.0 (5.3-28.7)
ACE inhibitors (n=14)	4.0 (1.9-6.1)	-	6.1 (0.0-17.0)	-
NSAID (n=16)	3.1 (1.6-4.5)	-	-	-
Oral Steroids (n=5)	1.3 (0.0-2.6)	-	-	2.6 (0.0-7.7)
Inhalator steroids (n=16)	4.8 (2.5-7.0)	-	-	3.7 (0.0-10.6)
Chronic diseases:				
High blood pressure (%) (n=76)	15.4 (12.1-18.7)	9.8 (28.8-16.8)	17.8 (5.5-30.1)	13.7 (4.1-23.2)
Diabetes (%) (n=31)	8.4 (5.6-11.1)	-	5.9 (0.0-16.0)	7.1 (0.0-15.8)
Asthma/COPD (n=33)	8.7 (5.6-11.8)	1.0 (0.0-3.1)	-	5.3 (0.0-12.4)
Cardiovascular disease (n=9)	1.7 (0.7-2.7)	-	-	-
Cancer (n=6)	1.1 (0.2-2.1)	-	-	-
Prevalence PCR-RT positive (n=100)	27.5 (22.1-32.8)	7.9 (3.4-12.3)	8.8 (0.0-18.4)	7.5 (0.0-15.6)
Prevalence of Ig M(+) or IgG (+) ever (n=70)	18.5 (14.1-23.0)	4.4 (0.5-8.2)	3.2 (0.0-9.4)	7.8 (0.0-16.2)
IgM (+) or IgG (+) by days of initial symptoms*:				
Baseline (Median: 3 days; (0-21)) (n=25)	5.7 (3.1-8.3)	3.3 (0.0-6.8)	3.9 (0.0-11.3)	2.7 (0.0-7.7)
Follow-up 1 (Median: 11 days; (6-24)) (n=7)#	1.9 (0.0-3.9)	1.0 (0.0-3.1)	2.5 (0.0-7.7)	3.0 (0.0-8.9)
Follow-up 2 (Median: 19 days (12-33)) (n=48)#	18.0 (13.3-22.6)	-	-	5.6 (0.0-13.2)
Initial symptoms profile (only symptomatic patients) :				
Symptoms	Prevalence % (95% CI)	Symptoms	Prevalence % (95% CI)	
Cephalaea (n=202)	66.9 (61.7-72.1)	Fatigue (n=49)	16.2 (12.1-20.4)	
Dry cough (n=141)	46.7 (41.1-52.3)	Productive cough (n=44)	14.6 (10.7-18.5)	
Sore throat (n=137)	45.4 (39.8-50.9)	Fever (n=44)	14.6 (10.6-18.5)	
Myalgia (n= 128)	42.4 (36.8-47.9)	Nasal congestion (n=42)	13.9 (10.0-17.8)	
Feverish sensation (n=83)	27.5 (22.5-32.5)	Chest pain (n=40)	13.2 (9.4-17.0)	
Diarrhea (n= 57)	18.9 (14.5-23.2)	Anosmia (n=30)	9.9 (6.6-13.3)	
Disnea (n=53)	17.5 (13.4-21.7)	Dysgeusia (n=22)	7.3 (4.4-10.2)	

Prevalence are adjusted by age, and sex using logistic regression. Age mean is adjusted by sex using linear regression

follow-up 1: follow-up in first time period N = 381

follow-up 2: follow-up in second time period N = 447

Table 2. Relative sensitivity, specificity and likelihood ratios associated with serologic tests compared with RT-PCR test results in symptomatic patients by time period.

Symptomatic patients, <=7 days since first symptom					
	RT-PCR (+)	RT-PCR (-)	Total	Sensitivity	4.9% (1.0-13.7%)
IgM or IgG (+)	3	5	8	Specificity	96.9% (92.9-99.0%)
IgM or IgG r (-)	58	157	215	LR+	1.59 (0.39-6.47)
Total	61	162	223	LR-	0.98 (0.92-1.05)
Symptomatic patients, 8 to 14 days					
	RT-PCR (+)	RT-PCR (-)	Total	Sensitivity	27.3% (6.0-61.0%)
IgM or IgG (+)	3	2	5	Specificity	98.7% (95.4-99.8%)
IgM or IgG r (-)	8	154	162	LR+	21.27 (3.96-114.3)
Total	11	156	167	LR-	0.74 (0.51-1.06)
Symptomatic patients >=15 days					
	RT-PCR (+)	RT-PCR (-)	Total	Sensitivity	57.8% (44.8-70.1%)
IgM or IgG (+)	37	3	40	Specificity	98.4% (95.5-99.7%)
IgM or IgG r (-)	27	187	214	LR+	36.6 (11.69-114.71)
Total	64	190	254	LR-	0.43 (0.32-0.57)

RT-PCR: Reverse transcriptase polymerase chain reaction
 LR: likelihood ratio

Figure 1. Positive rate (%) of IgM or IgG in symptomatic and asymptomatic patients with positive RT-PCR over time.

