

Table S1 – (Un)provided CV-care program per risk categories (n=391)

	Total	CV-care program primary care	CV-care program secondary care	No CV-care program
	n	n (%)	n (%)	n (%)
Very high CVR	178	127 (71.3)	9 (5.1)	42 (23.6)
DM	73	66 (90.4)	5 (6.8)	2 (2.7)
CVD	84	53 (63.1)	4 (4.9)	27 (32.1)
CKD	4	3 (75.0)	0 (0.0)	1 (25.0)
SCORE > 10%	2	0 (0.0)	0 (0.0)	2 (100)
> 70 years*	15	5 (33.3)	0 (0.0)	10 (66.7)
High CVR	119	81 (68.1)	1 (0.8)	37 (31.1)
DM	14	13 (92.9)	1 (7.1)	0 (0.0)
CKD	6	4 (66.7)	0 (0.0)	2 (33.3)
Severe RF	2	0 (0.0)	0 (0.0)	2 (100)
SCORE 5 – 9%	12	3 (25.0)	0 (0.0)	9 (75.0)
CV-medication	85	61 (71.8)	0 (0.0)	24 (28.2)
Low to mod. CVR	32	7 (21.9)	0 (0.0)	25 (78.1)
SCORE < 5%	31	7 (22.6)	0 (0.0)	24 (77.4)
< 40 years*	1	0 (0.0)	0 (0.0)	1 (100)
Unknown CVR	62	0 (0.0)	0 (0.0)	62 (100)

*with complete data on SCORE

n number, CV cardiovascular, CVD cardiovascular diseases, DM diabetes mellitus, CKD chronic kidney disease, RF risk factor, SCORE systemic coronary risk evaluation

Table S2 – Characteristics of patients with unknown CVR based on incomplete cardiovascular risk measurements

	Unknown risks, <i>n</i> = 62	
	n	%
Age < 40 years	0	0.0
Age 40 - 70 years	52	83.9
Age > 70 years	10	16.1
Smoking status		
Current	30	48.4
Former	17	27.4
Never	5	8.1
<i>Missing</i>	10	16.1
Systolic BP > 140 mmHg	11	17.7
<i>Missing</i>	16	25.8
LDL-c > 2.5 mmol/L	3	4.8
<i>Missing</i>	58	93.6
BMI > 25 kg/m ²	19	29.2
<i>Missing</i>	23	37.1
eGFR < 60 mL/min/1.73m ²	2	3.2
<i>Missing</i>	41	66.1
ACR-u (mg/mmol)	1.0	
<i>Missing</i>	61	98.4
Patients characteristics ≥ 60 years (n=29)		
Frailty		
Yes	12	41.4
No	9	31.0
<i>Missing</i>	8	27.6
Polypharmacy		
Yes	7	24.1
No	13	44.8
<i>Missing</i>	9	31.0

n number, BP blood pressure, LDL-c low-density lipoprotein-cholesterol,

BMI body mass index, eGFR estimated glomerular filtration rate,

ACR albumin-creatinine ratio in urine